

Acceptance and commitment therapy for psychosis: randomised controlled trial

Frances Shawyer, John Farhall, Neil Thomas, Steven C. Hayes, Robert Gallop, David Copolov and David J. Castle

Background

The efficacy of acceptance and commitment therapy (ACT) in psychosis has been reported but not for medication-resistant psychosis.

Aims

To test the efficacy of ACT in a sample of community-residing patients with persisting psychotic symptoms. (Australian New Zealand Clinical Trials Registry: ACTRN12608000210370.)

Method

The primary outcome was overall mental state at post-therapy (Positive and Negative Syndrome Scale – total); secondary outcomes were psychotic symptom dimensions and functioning. In total, 96 patients were randomised to ACT ($n=49$) or befriending ($n=47$). Symptom, functioning and process measures were administered at baseline, post-therapy and 6 months later.

Results

There was no group difference on overall mental state. In secondary analyses the ACT group showed greater improvement in positive symptoms and hallucination distress at follow-up: Cohen's $d=0.52$ (95% CI 0.07–0.98) and 0.65 (95% CI 0.24–1.06), respectively.

Conclusions

Improvements reflected the treatment focus on positive symptoms; however, absence of process-measure changes suggests that the ACT intervention used did not manipulate targeted processes beyond befriending. Symptom-specific therapy refinements, improved investigation of process and attention to cognitive functioning and dose are warranted in future research.

Declaration of interest

None.

Copyright and usage

© The Royal College of Psychiatrists 2016.

The prevalence of medication-resistant psychotic symptoms has led to psychological treatment being used alongside medication to reduce the impact of symptoms. Cognitive-behavioural therapy (CBT) for psychosis (CBTp) has been the dominant approach, but is lengthy and complex to deliver.¹ Substantial interest has grown in applying acceptance and mindfulness-based therapies,² with acceptance and commitment therapy (ACT) – one of the most widely used manualised approaches^{3,4} – showing promise. Rather than effecting change by modifying the person's understanding of psychotic experiences, ACT targets the extent to which symptoms and related beliefs dominate conscious experience and behaviour.⁵ Two randomised controlled trials (RCTs) have tested four-session ACT interventions in in-patients with acute psychosis, observing small-to-medium reductions in re-admission to hospital rates over 4–12 months in intention-to-treat analyses *v.* routine or enhanced routine care.^{6–8} However, ACT has yet to be trialled for medication-resistant psychosis. The aim of this study was to test the efficacy of ACT in a sample of community-residing patients with persisting psychotic symptoms. This paper reports our results for primary and secondary outcome measures. Additionally, several process measures were included in order to examine hypothesised mechanisms of action (see study protocol⁹). Our hypotheses were that, compared with patients receiving equivalent clinician time in a comparison condition, patients who received ACT would show improvements in (a) overall mental state; (b) preoccupation, conviction, distress and disruption to life associated with positive symptoms; and (c) social functioning. We hypothesised that these changes would be achieved by the conclusion of therapy and maintained at 6-month follow-up. Outcomes were also examined in relation to service utilisation.

Method

Design

The study protocol, detailed in Thomas *et al.*,⁹ is briefly described here. A prospective single (rater)-blind RCT compared two parallel groups: the intervention (ACT) *v.* a comparison condition (befriending). Assessments took place prior to randomisation (baseline), at post-therapy and 6 months after the end of therapy. The trial is registered at the Australian New Zealand Clinical Trials Registry: ACTRN12608000210370.

Participants

Recruitment took place over 3 years from public community mental health services, non-government psychiatric rehabilitation services and private providers in Melbourne, Australia, supplemented by media advertising. Inclusion criteria were: (a) aged 18–65 years; (b) current diagnosis of schizophrenia or schizoaffective disorder; (c) residual hallucinations or delusions associated with significant distress or disability (score ≥ 4 on the Positive and Negative Syndrome Scale (PANSS)¹⁰ items P1 and/or P3); (d) these symptoms present continuously over the past 6 months; and (e) on therapeutic doses of antipsychotic medication over the past 6 months (clinician report). Exclusion criteria were: (a) any neurological disorder that may affect cognitive function; (b) insufficient conversational English; (c) Wechsler Test of Adult Reading (WTAR)¹¹ estimated IQ < 70 ; (d) change of antipsychotic medication within the previous 8 weeks or planned at the time of intake; (e) currently receiving other formal psychological treatment. Following consent,

participants were interviewed to confirm eligibility and complete baseline assessment measures prior to randomisation.

Sample size

The target sample size of 53 participants per treatment arm was calculated to detect post-therapy between-group effects for overall mental state of $d = 0.55$ or greater with 80% power ($\alpha = 0.05$). This effect size was slightly lower than the $d = 0.60$ reported in the TORCH¹² and Gaudiano & Herbert⁸ trials for overall mental state, in view of the novel protocol.

Treatment conditions

ACT

Participants were offered eight 50 min ACT sessions, delivered weekly to fortnightly over around 3 months. ACT was conducted according to a local manual based on the (transdiagnostic) ACT manual³ with recommended adaptations for psychosis.^{6,13,14} Participants were provided with handouts and sessions recorded onto compact discs for home review.

Befriending

Participants were offered eight 50 min sessions of the befriending intervention,¹⁵ a manualised treatment previously used as a control condition in psychological intervention trials in psychosis,^{12,16,17} befriending involves engaging in conversation about everyday topics, while overtly avoiding discussion of symptoms and problems. It produces similar treatment expectancy and engagement to CBT, with similar drop-out rates.¹⁸ Befriending has shown some equivalence in outcomes to CBTp, suggesting that it has effectiveness as a treatment in its own right.^{12,16,17}

In both conditions, therapy was provided by four clinical psychologists, experienced in psychological interventions for schizophrenia, with additional training in ACT and befriending. They attended weekly peer supervision led by J.F. or S.C.H. Local services managed medication, case management and other aspects of treatment.

Treatment fidelity

An independent assessor, masked to treatment allocation, rated a stratified random sample of treatment session audio files for adherence with each therapy protocol, and assigned each session to ACT or befriending. The Befriending Treatment Integrity Measure (BTIM)¹⁵ was used to assess the quality of befriending sessions and to ensure that ACT sessions did not include befriending techniques. In the absence of a suitable ACT fidelity scale, we developed the six-item ACT for Psychosis Adherence and Competence Scale (APACS – see online supplement DS1), where adherence ratings reflect a composite of the presence and frequency of the six ACT processes defined in our manual. The APACS adherence subscale showed acceptable psychometric properties.¹⁹

Assessments

Research assistants masked to treatment condition administered the eligibility and assessment measures in face-to-face interviews with participants. The Structured Clinical Interview for DSM-IV-TR Axis I Disorders²⁰ and WTAR were completed at baseline to assess eligibility. Symptom outcome measures included the following: PANSS, assessing overall mental state including positive, negative and general symptoms; the Psychotic Symptom Rating Scales (PSYRATS),²¹ assessing the frequency, preoccupation, conviction, distress and disruption to life associated with auditory hallucinations (PSYRATS-AH) and main delusions (PSYRATS-D); and the Peters Delusions Inventory

(PDI)²² to assess the range/number of delusional beliefs held and overall degree of associated distress, preoccupation and conviction. The PSYRATS-AH do not include a specific item assessing preoccupation so we created an additional item assessing time spent thinking about voices. Psychosocial functioning was measured by the Social Functioning Scale (SFS).²³ Service utilisation related to psychiatric hospital admissions and mental health consultations was also assessed.⁹

Process measures included the 16-item Acceptance and Action Questionnaire (AAQ)²⁴ to assess (a) acceptance of psychotic experiences as opposed to experiential avoidance, and (b) commitment to valued action; the Voices Acceptance and Action Scale (VAAS)²⁵ was used to assess acceptance and autonomous action in relation to auditory hallucinations; and the Recovery Style Questionnaire (RSQ)²⁶ to assess the degree to which participants 'integrate' their illness, acknowledging their illness experiences with interest, as opposed to 'sealing over', seeking to separate psychosis from themselves. Additional measures were administered⁹ but are not reported here.

Measures, and dose of antipsychotic medication, were assessed at each time point except for service utilisation, administered at baseline and follow-up. At the end of post-therapy assessments, participants were asked to rate therapy acceptability on the Client Satisfaction Questionnaire (CSQ)²⁷ (possible scores: 8–32). Additional therapy evaluation questions assessed emotional response to sessions and extent of improvement of problems related to psychosis. Rater's masking was preserved by participants returning responses in a sealed envelope. Baseline assessments commenced in October 2008 with the final follow-up conducted in November 2012.

Randomisation and masking

Randomisation⁹ was prepared by an independent statistician. Stratification was by site and recovery style (integration or sealing over) giving 18 factorial groups. Allocation was by a random permuted blocks procedure within Microsoft Excel, using a random number generator to choose each sequence of blocks of sizes 2, 4 and 6, without replacement. Numbered, sealed, opaque envelopes for each of the 18 groups' concealed allocations. Using the envelopes in order, for the appropriate group, an independent researcher allocated participants to treatment with a 1:1 allocation ratio in accordance with CONSORT guidelines.

Considerable efforts were made to maintain rater masking.⁹ Masking was assessed by asking raters to classify participants into a treatment condition after post-therapy and follow-up assessments and indicate their level of confidence. Breaches in masking were recorded and addressed by changing the rater wherever possible.

Primary and secondary outcomes

The primary outcome is PANSS total at post-therapy, with PANSS positive, negative and general subscales also reported. Secondary outcomes include PSYRATS-AH and PSYRATS-D preoccupation, conviction, distress and disruption to life, adjusting for the presence and frequency of symptoms; the SFS; and service utilisation.

Statistical analysis

The analyses were not masked. Where distributions were non-normal across time points, square-root transformations were applied; descriptive statistics are reported for back-transformed data. As a result of some outcome measures showing significant baseline differences between ACT and befriending, and between completers and non-completers, propensity score matching²⁸ was used to even out the distribution of the measured baseline

characteristics across the intervention groups. This involved constructing a logistic regression model with treatment condition as the outcome, and baseline clinical variables and demographics as predictors, plus a 'missing-at-post-therapy' variable to index differential effects of attrition. Based on this model, the probability of being chosen for the 'reference' group (propensity score) was estimated for each participant, to provide a summary of the covariate imbalance between intervention and control participants. These propensity scores were included as a covariate in all analyses (excluding therapy evaluation measures).²⁹ Intention-to-treat analyses using mixed regression models were the primary analytic approach. Compared with other techniques such as repeated measures analysis of variance, the mixed-model framework deletes randomly missing observations without dropping the participant. All participants randomised are included in the analysis; therefore, imputation methods such as last observation carried forward (LOCF) or expectation-maximisation are not applied. The within-groups factor was time (baseline, post-therapy, follow-up) and the between-groups factor was group (ACT, befriending). For service utilisation, the within-groups factor was time (baseline, follow-up). Reports are based on the covariance model with the fewest parameters that were not significantly different than the most complex model as determined by comparison of nested models through the restricted log-likelihood. Both mixed model repeated measures (MMRM), which treats time as categorical, and hierarchical linear modelling (HLM), which treats time as a linear covariate, were calculated and compared. In all cases MMRM provided a better fit to the data as determined by a comparison of the restricted log-likelihood, and thus MMRM results are reported throughout as the primary analysis. Because

HLM better addresses overall trends across time, these analyses are reported if they conveyed additional information.

The PSYRATS analyses were conducted on individual items using ordinal regression models³⁰ to accommodate both within- and between-participant clustering. Analyses for PSYRATS-AH and PSYRATS-D scales included only data from participants reporting auditory verbal hallucinations or delusions, respectively, during the project. Analyses related to the PSYRATS-AH subscales included PSYRATS-AH frequency as a covariate. Analyses for the PSYRATS-D subscales included PSYRATS-D amount of preoccupation (frequency of delusional thinking) as a covariate.

Planned contrasts were used to compare changes from baseline under each intervention at post-therapy and follow-up. Chi-square tests were used to compare the number of participants in the ACT and befriending groups who achieved a clinically significant improvement on PANSS scores, defined as a 25% score reduction from baseline as recommended by Leucht.³¹ Analyses were undertaken using SPSS 21 for Windows: the linear mixed-effects models (MIXED) procedure was used for the MMRM analyses. The ordinal regression model for the PSYRATS data was implemented in SAS 9.4 with the SAS Procedure PROC GLIMMIX. GLIMMIX was used to predict the probability of being in the lower category.

Results

Participant characteristics

Participants included 59 males (61.5%) and 37 females (38.5%), mean age: 36.1 years (s.d.=9.1, range 19–64). Baseline clinical and demographic characteristics are shown in Table 1.

Variable	ACT group (n = 49)	Befriending group (n = 47)
Age, years: mean (IQR)	35.6 (15.3)	33.0 (8.5)
Gender, n (%)		
Men	29 (59.2)	30 (63.8)
Women	20 (40.8)	17 (36.2)
Marital status, n (%)		
Single	36 (73.5)	39 (83.0)
Married/de facto	6 (12.2)	5 (10.6)
Divorced/separated/widowed	7 (14.3)	3 (6.4)
Education status, n (%) ^b		
Secondary	37 (77.1)	30 (63.8)
Certificate/diploma	8 (16.7)	11 (23.4)
Tertiary	3 (6.3)	6 (12.8)
Main occupation past 7 days, n (%)		
Employed (part-time/casual)	7 (14.3)	7 (14.9)
Volunteer	3 (6.1)	2 (4.3)
Student (part time)	3 (6.1)	2 (4.3)
Home duties/retired	2 (4.1)	1 (2.1)
Unemployed	34 (69.4)	35 (74.5)
Disability Support Pension ^c	42 (87.5)	43 (95.6)
Wechsler Test of Adult Reading IQ, mean (s.d.) ^c	99.9 (8.7)	101.4 (8.5)
DSM-IV diagnosis, n (%)		
Schizophrenia	35 (71.4)	38 (80.9)
Schizoaffective disorder	14 (28.6)	9 (19.1)
Positive and Negative Syndrome Scale score		
Positive subscale, mean (s.d.)	22.8 (5.5)	20.8 (5.2)
Negative subscale, mean (s.d.)	16.9 (4.4)	19.2 (5.3)
General subscale, mean (s.d.)	40.0 (12.0)	37.0 (9.0)
Total score, median (IQR)	77.0 (23.0)	75.0 (21.0)
Chlorpromazine equivalent dose, mg: median (IQR)	778.8 (462.0)	840.0 (507.0)

a. Medians (interquartile range (IQR)) reported where data are skewed.
b. n = 48 for the ACT group and n = 47 for the befriending group.
c. n = 48 for the ACT group and n = 45 for the befriending group.

Participant flow, attrition and reliability

Of 531 patients identified as possible candidates, 137 either declined participation before eligibility could be assessed or were unable to be assessed. The eligibility of 244 of the remaining 394 patients was confirmed, however 83 were unavailable for participation. Of the 161 eligible and available, 65 declined participation and 96 (60% of eligible candidates) proceeded to randomisation (Fig. 1). There were no significant group differences in rates of attrition (post-therapy: $\chi^2(1)=0.06$, $P=0.81$; follow-up: $\chi^2(1)=0.76$, $P=0.38$). Completers ($n=77$) at follow-up were compared with participants who did not

complete the final assessment ($n=19$) on baseline measures. Non-completers showed more frequent delusion-related distress (PSYRATS-D amount of distress), and lower acceptance (AAQ, VAAS-9). PANSS interrater reliability assessment included all participants with a second rating ($n=23$). Median intraclass correlations ranged from 0.97 (PANSS-Total) to 0.84 (PANSS-General).

Fidelity

A masked fidelity assessor correctly assigned to condition all 94 sampled sessions. The mean total APACS score across the 48




Fig. 1 Flow of participants through the study (CONSORT diagram).

a. Includes unstable mental state, unstable circumstances, discharge from service, move from area, risk issues, unreliable attendance and poor engagement with services. ACT, Acceptance and Commitment Therapy; ITT, intention to treat.

ACT sessions sampled was 7.06 (s.d. = 1.79) indicating, on average, that the majority of each session involved identifiable ACT processes. One sampled befriending session had a rated ACT component (values); no ACT sessions had a befriending component rated.

Masking

Only one assessment (at follow-up) was conducted unmasked during the trial (because no other rater was available). Data on raters' guess of treatment group were available for 33 interviews at post-therapy and 50 at follow-up. Using one-tailed binomial tests, neither post-therapy nor follow-up guesses were better than chance. Mean confidence ratings were 14.6 (s.d. = 18.7) at post-therapy and 17.8 (s.d. = 25.3) at follow-up, and were unrelated to accuracy.

Treatment exposure

Participants allocated to ACT completed an average of 7.0 (s.d. = 2.3) therapy sessions, compared with 7.2 (s.d. = 2.0) for befriending, Mann-Whitney *U*-test, *P* = 0.82. There was no group difference in duration of therapy (months) (ACT group 2.9, s.d. = 1.3; befriending group 2.6 (s.d. = 0.8), Mann-Whitney *U*-test, *P* = 0.36). There were no significant changes by group in chlorpromazine-equivalent antipsychotic medication dosages over the study period (time: *F*(2, 145.4) = 1.43, *P* = 0.24; group: *F*(1, 93.9) = 0.30, *P* = 0.59; group × time: *F*(2, 145.54) = 1.21, *P* = 0.30).

Main findings

Results of the MMRM and ordinal regression analyses for the key symptom, functioning and service utilisation outcomes are shown in Tables 2–4.

Primary outcomes

There were no significant group × time differences on the primary outcome measure, PANSS total (Table 2). The group × time result for PANSS positive showed an effect size of *d* = 0.37, which fell just short of significance. Further examination, with planned contrasts indicated that the participants in the ACT group had a significantly greater and medium improvement compared with the befriending group at follow-up, *t*(79.4) = -2.33, *P* = 0.02, *d* = 0.52 (95% CI 0.07–0.98) but not at post-therapy (*P* = 0.30). HLM analyses showed a statistically significant difference in slopes of improvement between conditions (group × time *F* = 5.59, d.f. = 84.35, *P* = 0.02, effect size 0.48 (95% CI 0.06–0.89)).³² Including IQ scores from the WTAR as a covariate strengthened findings also (group × time result for MMRM analysis: *F* = 3.30, d.f. = 78.25, *P* = 0.04, effect size 0.41 (95% CI 0.07–0.74); ACT *v.* befriending planned contrast at follow-up: *t*(76.6) = -2.85, *P* = 0.01, *d* = 0.59 (95% CI 0.17–1.00)).

Secondary outcomes

There were significant group × time interactions for PSYRATS-AH amount of distress (the amount of time voices are distressing) and PSYRATS-AH disruption to life (Table 3). Planned baseline to follow-up contrasts indicated that, compared with the befriending group, the participants in the ACT group reported significantly less voice-related distress, with a medium effect size, *t*(101) = -3.25, *P* = 0.002, *d* = 0.65 (95% CI 0.24–1.06). The voice-related disruption to life contrast showed a small effect size and did not reach significance, *t*(99) = 1.75, *P* = 0.08, *d* = 0.35 (95% CI -0.05 to 0.75). There were no significant group × time interactions for the remaining PSYRATS subscales or for the SFS

Table 2 Means (95% CI), Cohen's *d* values and tests of group × time fixed effects in mixed-effects model repeated measures ANOVA for Positive and Negative Syndrome Scale (PANSS) primary outcome measures^a

Measures	ACT group				Befriending group				Group × time
	Mean (95% CI)		Within group, <i>d</i> (95% CI), <i>P</i>		Mean (95% CI)		Within group, <i>d</i> (95% CI), <i>P</i>		
	Post-therapy	Follow-up	Baseline to post-therapy	Baseline to follow-up	Post-therapy	Follow-up	Baseline to post-therapy	Baseline to follow-up	
PANSS total ^b	74.2 (69.7 to 78.8)	72.4 (67.6 to 77.0)	0.33 (0.02 to 0.64)	0.46 (0.15 to 0.77)	71.7 (67.2 to 76.3)	73.3 (68.6 to 78.2)	0.42 (0.11 to 0.73)	0.30 (-0.24 to 0.56)	0.63 (164.6) 0.54 (-0.24 to 0.56)
PANSS positive	20.5 (18.8 to 22.1)	19.2 (17.5 to 21.0)	0.64 (0.32 to 0.96)	0.79 (0.47 to 1.11)	19.2 (17.5 to 20.9)	19.7 (17.8 to 21.6)	0.42 (0.11 to 0.73)	0.23 (-0.08 to 0.54)	2.82 (81.2) 0.07 (-0.03 to 0.77)
PANSS negative	16.9 (15.5 to 18.4)	17.4 (15.9 to 18.9)	-0.01 (-0.33 to 0.33)	-0.06 (-0.36 to 0.24)	17.6 (16.1 to 19.1)	17.6 (16.1 to 19.2)	0.25 (-0.06 to 0.56)	0.24 (-0.07 to 0.55)	1.14 (166.0) 0.32 (-0.09 to 0.53)
PANSS general ^b	37.0 (34.5 to 39.5)	35.9 (33.4 to 38.4)	0.27 (-0.04 to 0.58)	0.41 (0.10 to 0.72)	35.1 (32.6 to 37.6)	36.0 (33.4 to 38.7)	0.37 (0.06 to 0.68)	0.26 (-0.05 to 0.57)	0.65 (165.1) 0.52 (-0.24 to 0.56)

ACT, acceptance and commitment therapy.
 a. Propensity scores included as a covariate. Within-group negative *d* value indicates deterioration.
 b. Analysis based on square root transformed data to correct positive skew, back-transformed means and confidence intervals are shown.

Table 3 Means (95% CI), Cohen's *d* values and group × time fixed effects in ordinal regression analyses for Psychotic Symptom Rating Scales (PSYRATS) secondary outcome measures^a

Measures	ACT group						Befriending group						Group × time			
	Mean (95% CI)		Follow-up		Within group, <i>d</i> (95% CI), <i>P</i>		Mean (95% CI)		Follow-up		Within group, <i>d</i> (95% CI), <i>P</i>		F (d.f.) <i>P</i>		<i>d</i> (95% CI)	
	Baseline	Post-therapy	Follow-up	Baseline to post-therapy	Baseline to follow-up	Baseline	Post-therapy	Follow-up	Baseline to post-therapy	Baseline to follow-up	F	(d.f.)	<i>P</i>	<i>d</i>	(95% CI)	
PSYRATS-AH^b																
Preoccupation	2.44 (1.99 to 2.89)	2.32 (1.96 to 2.68)	2.27 (1.73 to 2.82)	0.11 (-0.29 to 0.50)	0.02 (-0.37 to 0.42) 0.91	2.21 (1.65 to 2.77)	1.93 (1.3 to 2.56)	2.29 (1.6 to 2.98)	0.09 (-0.31 to 0.48) 0.67	0.38 (0.02 to 0.78) 0.06	1.22	(100)	0.30	0.23	(-0.17 to 0.63)	
Origin beliefs	2.51 (2.1 to 2.92)	2.23 (1.75 to 2.71)	2.62 (2.1 to 3.14)	0.11 (-0.21 to 0.43) 0.49	-0.12 (-0.20 to 0.44) 0.45	2.66 (2.17 to 3.15)	2.44 (1.86 to 3.03)	2.30 (1.58 to 3.02)	0.02 (-0.30 to 0.34) 0.90	0.06 (-0.26 to 0.38) 0.69	0.66	(152)	0.52	0.17	(-0.15 to 0.49)	
Amount of distress	2.72 (2.37 to 3.07)	2.23 (1.78 to 2.68)	2.10 (1.59 to 2.61)	0.33 (-0.07 to 0.73) 0.10	0.43 (0.03 to 0.83) 0.03	2.31 (1.78 to 2.84)	2.22 (1.61 to 2.83)	2.57 (1.83 to 3.31)	0.10 (-0.29 to 0.49) 0.61	0.49 (0.09 to 0.89) 0.02	5.27	(101)	0.01	0.47	(0.07 to 0.87)	
Intensity of distress	2.64 (2.32 to 2.97)	2.32 (1.89 to 2.75)	2.07 (1.54 to 2.6)	0.18 (-0.21 to 0.58) 0.36	0.33 (-0.07 to 0.72) 0.11	2.59 (2.06 to 3.12)	2.11 (1.53 to 2.7)	2.29 (1.57 to 3.01)	0.20 (-0.19 to 0.60) 0.31	0.06 (-0.33 to 0.45) 0.75	1.18	(101)	0.31	0.22	(-0.18 to 0.62)	
Disruption of life	1.89 (1.69 to 2.09)	1.66 (1.37 to 1.95)	1.30 (0.95 to 1.66)	0.21 (-0.19 to 0.61) 0.30	0.58 (0.17 to 0.99) 0.005	1.66 (1.3 to 2.02)	1.15 (0.77 to 1.53)	1.43 (0.94 to 1.92)	0.43 (0.03 to 0.84) 0.03	0.04 (-0.35 to 0.44) 0.83	3.27	(99)	0.04	0.37	(0.00 to 0.74)	
PSYRATS-D^c																
Preoccupation	2.51 (2.12 to 2.91)	1.97 (1.46 to 2.48)	1.90 (1.43 to 2.37)	0.33 (-0.08 to 0.74) 0.11	0.42 (0.00 to 0.83) 0.045	2.62 (2.1 to 3.15)	2.00 (1.52 to 2.48)	2.10 (1.53 to 2.67)	0.14 (-0.27 to 0.54) 0.51	0.06 (-0.34 to 0.46) 0.77	0.58	(95)	0.56	0.16	(-0.20 to 0.52)	
Conviction	2.85 (2.39 to 3.31)	2.32 (1.79 to 2.85)	2.21 (1.61 to 2.81)	0.36 (-0.05 to 0.78) 0.08	0.39 (-0.03 to 0.80) 0.06	3.27 (2.81 to 3.73)	2.70 (2.15 to 3.25)	2.85 (2.17 to 3.53)	0.25 (-0.16 to 0.66) 0.23	0.07 (-0.34 to 0.48) 0.74	0.45	(94)	0.64	0.14	(-0.22 to 0.50)	
Amount of distress	2.71 (2.29 to 3.13)	2.19 (1.60 to 2.78)	1.90 (1.35 to 2.45)	0.16 (-0.16 to 0.49) 0.32	0.41 (0.08 to 0.74) 0.01	2.69 (2.16 to 3.22)	2.11 (1.5 to 2.72)	2.1 (1.36 to 2.84)	0.09 (-0.23 to 0.41) 0.59	0.07 (-0.25 to 0.39) 0.68	0.87	(150)	0.42	0.19	(-0.18 to 0.56)	
Intensity of distress	2.74 (2.36 to 3.13)	1.84 (1.33 to 2.35)	1.79 (1.24 to 2.34)	0.57 (0.16 to 0.99) 0.006	0.65 (0.23 to 1.06) 0.002	2.73 (2.24 to 3.22)	1.93 (1.36 to 2.5)	1.80 (1.16 to 2.44)	0.31 (-0.10 to 0.72) 0.14	0.45 (0.03 to 0.86) 0.03	0.29	(95)	0.75	0.11	(-0.25 to 0.47)	
Disruption of life	1.76 (1.53 to 1.99)	1.17 (0.86 to 1.48)	1.34 (0.97 to 1.71)	0.61 (0.19 to 1.03) 0.004	0.45 (0.03 to 0.87) 0.03	1.65 (1.25 to 2.05)	1.07 (0.69 to 1.45)	1.37 (0.91 to 1.83)	0.44 (0.02 to 0.85) 0.04	0.11 (-0.30 to 0.51) 0.61	0.50	(93)	0.61	0.15	(-0.21 to 0.51)	

ACT, acceptance and commitment therapy.
a. Propensity scores included as a covariate. Italicised *d* value indicates deterioration.
b. PSYRATS auditory hallucinations (PSYRATS-AH) frequency included as a covariate.
c. PSYRATS delusions (PSYRATS-D) amount of preoccupation included as a covariate.

(Table 4). Although the study was not powered to assess changes to rates of hospital admission,⁹ these appeared similar at follow-up to baseline in both groups. (See online Table DS3 for untransformed means and standard deviations for primary and secondary measures.)

Other outcomes. On the remaining outcome measures, only the PDI distress subscale showed a significant group × time interaction ($F(151.3) = 3.46, P = 0.03, d = 0.38$), with the befriending group reporting a medium and significantly lower level of delusion-related distress at follow-up, $t(155.9) = 2.62, P = 0.01, d = 0.62$, but not at post-therapy ($P = 0.32$).

Clinical significance. At follow-up, 51.2% (21/41) of the ACT group showed a clinically significant reduction in positive symptoms (i.e. a 25% reduction in PANSS positive), compared with only 22.2% (8/36) of the befriending group ($\chi^2(1, n = 77) = 6.89, P = 0.009, d = 0.63$). There were no significant group differences on PANSS total or the other PANSS subscales (see online Table DS4).

Process measures. There were no significant group × time interactions for any of these measures (see online Table DS5) precluding exploration of mechanisms.

Therapy evaluation. Based on data from the CSQ, mean levels of satisfaction with therapy were significantly higher in the ACT group (mean 26.6, s.d. = 3.9, $n = 40$) compared with the befriending group (mean 23.9, s.d. = 4.5, $n = 41$), with a medium effect size; $t(79) = -2.97, P = 0.004, d = 0.65$. The ACT group also endorsed significantly higher levels of problem improvement (76.9%, $n = 39$ v. 43.2%, $n = 37, \chi^2 = 9.0, P = 0.003, d = 0.73$) and emotional improvement (89.7%, $n = 39$ v. 59.5%, $n = 37, \chi^2 = 9.3, P = 0.002, d = 0.75$), both with medium effect sizes.

Discussion

Compared with the befriending group, the participants in the ACT group were more satisfied with therapy and reported greater subjective benefit. Hypothesised greater gains for the ACT group on masked ratings were, however, only partially supported. There were no group differences on PANSS total, and the difference on PANSS positive using MMRM analyses fell just short of significance. Despite not meeting this convention, several findings are suggestive of a relationship favouring the ACT group for PANSS positive: the relationship was significant using HLM analyses and with MMRM when IQ scores were added as a covariate, and the original planned comparison estimated a medium effect size in favour of ACT at follow-up. Further, around half the ACT group achieved clinically significant improvement on PANSS positive at follow-up compared with less than a quarter of the befriending group ($P < 0.05$). These findings were consistent with the ACT group showing significantly greater improvements at follow-up for PSYRATS-AH amount of distress. These changes correspond to the treatment focus on persisting positive symptoms, and the effect sizes appear comparable with the small-to-medium effects observed in similar trials of CBTp.^{33,34} However, caution is warranted because no significant group differences in favour of ACT were observed for the other outcome measures, and the befriending group unexpectedly showed significantly greater improvement in reported delusional distress, with a medium effect at follow-up. Additionally, no group differences were evident for any of the process measures.

Table 4 Means (95% CI), Cohen's *d* values and group × time fixed effects in mixed-effects model repeated measures ANOVA for social functioning and service utilisation^a

Measures	ACT group				Befriending group				Group × time			
	Mean (95% CI)		Within group, <i>d</i> (95% CI), <i>P</i>		Mean (95% CI)		Within group, <i>d</i> (95% CI), <i>P</i>					
	Baseline	Post-therapy	Follow-up	Baseline to post-therapy	Baseline to follow-up	Baseline	Post-therapy	Follow-up		Baseline to post-therapy	Baseline to follow-up	
Social Functioning Scale	100.4 (97.7 to 103.2)	99.7 (96.8 to 102.6)	99.5 (96.6 to 102.4)	-0.12 (-0.48 to 0.24) 0.43	-0.16 (-0.52 to 0.20) 0.32	99.6 (96.7 to 102.5)	100.2 (97.2 to 103.1)	99.7 (96.7 to 102.8)	0.09 (-0.27 to 0.45) 0.54	0.03 (-0.33 to 0.39) 0.88	0.57 (158.8) 0.57	0.15 (-0.21 to 0.51)
Service utilisation, number of consultations ^b	26.7 (21.1 to 33.0)	-	26.6 (20.5 to 33.5)	-	0.01 (-0.35 to 0.37) 0.96	27.4 (21.6 to 34.0)	-	19.5 (14.0 to 25.9)	-	0.40 (0.03 to 0.77) 0.02	2.82 (75.1) 0.097	0.39 (-0.07 to 0.85)
Service utilisation, length of consultations, ^b min	792.0 (618.8 to 986.5)	-	970.0 (672.4 to 1322.0)	-	-0.18 (-0.54 to 0.18) 0.27	775.5 (600.3 to 973.1)	-	713.1 (443.7 to 1046.1)	-	0.01 (-0.35 to 0.37) 0.69	1.12 (70.1) 0.29	0.25 (-0.22 to 0.73)

ACT, acceptance and commitment therapy.
 a. Propensity scores included as a covariate. Within-group negative *d* value indicates deterioration.
 b. Analysis based on square root transformed data to correct positive skew; back-transformed means and confidence intervals are shown.

Effect on positive symptoms

In considering the possibility of benefits specific to hallucinations, it is notable that the improvements in re-admission to hospital rate reported in the Bach & Hayes trial⁶ were stronger for participants with auditory hallucinations (ACT group 11.8%; treatment as usual 58.9%) than those with delusions (ACT group 38.5%, treatment as usual 28.5%). The Gaudio & Herbert trial⁸ also reported favourable findings for participants with auditory hallucinations (hallucination-related distress) but had too few self-ratings of delusions ($n = 8$) for analysis. It is possible that dose may be a factor here also with delusions potentially requiring longer therapy contact, as was suggested by Bach & Hayes.⁶ Although we considered that eight sessions, twice the number of Bach & Hayes, would provide a more comprehensive treatment for our chronically affected sample, this is substantially fewer than in trials of CBTp.³⁵ Indeed, compared with the acute phase, when normal coping has been disrupted, it may be more challenging to facilitate change with what may be well-developed and entrenched ways of adapting to symptoms during the chronic phase. Examining mean scores across the 14 symptom measures (Tables 2–3) shows a pattern of linear improvement over the three time points for nine measures in ACT but only two in befriending. In contrast, a pattern of improvement to post-therapy then loss of gain at follow-up is observed for ten measures in befriending but only two in ACT. Although speculative, it is possible that a longer follow-up period is needed to demonstrate clear group differences.

Results for delusions have also been less consistent using CBTp. A recent meta-analysis examining CBTp effects in auditory hallucinations and delusions³ found evidence of greater amenability to change of hallucinations *v.* delusions. If so, future studies may best examine more targeted methods and effects rather than examine effects on positive symptoms in combination.^{34,36–38}

It should also be noted that ACT protocols vary with problem presentation³⁹ and ACT for psychosis is still in its infancy thus further protocol refinement is likely required. For example, entrenched avoidant adaptation is a key challenge in chronic presentations: our preliminary recommendation is that dose be consistent with the 15 plus sessions established for CBTp and that the procedures on cost of current coping and engendering hope for change be extended. It has been suggested from an ACT perspective that hallucinations are often a target of experiential avoidance,⁶ whereas delusions are a means of avoidance – if adopting an ACT approach, each symptom may require different strategies. For example, given the observed reduction in delusional distress and reduced service use in the befriending group, an ACT protocol tailored to delusions might prioritise focus on valued living.

Processes of change

The fact that our ACT process measures did not change differentially in the ACT group leaves unclear the processes leading to improvements with hallucinations, making the current study more a test of a protocol than the underlying model. It suggests a need for additional protocol development given that psychological flexibility changes with ACT interventions in other populations, and in people with psychosis. The process of change is unlikely to be intrusion of cognitive strategies used in CBTp: session ratings using the Revised Cognitive Therapy for Psychosis Adherence Scale⁴⁰ suggested that very few cognitive techniques were used.¹⁹ It is also possible that more general processes, such as using a structured collaborative approach to respond adaptively to hallucinations, is an effective ingredient, aside from specific ACT interventions.

A qualitative study with a subset of ACT participants showed that although some participants were able to articulate an

understanding of ACT processes and attribute positive change to these, others found it difficult to understand exercises and were ambivalent about the usefulness of some aspects of ACT.⁴¹ In future work, it will be worthwhile identifying for whom ACT is most suited. For example, participants with cognitive impairments appeared to struggle with the specific ACT intervention tested in this trial. Adding WTAR scores as a covariate slightly improved outcomes for ACT in the area of positive symptoms suggesting, at least, a role for verbal learning in responsiveness to ACT as used in this study. Alternatively, simpler methods of teaching ACT concepts may need to be developed for this subpopulation.

Negative symptom change and functioning

Contrary to hypotheses, ACT had no significant impact on negative symptoms and functioning. This contrasts with the Gaudio & Herbert trial,⁸ which reported greater improvements in social functioning for ACT, and a pilot ($n = 27$) trial by White *et al*,⁴² who reported small improvements in negative symptoms following up to ten ACT sessions. These differences could be because of population, protocol, comparison condition or other factors.

Strengths and limitations

This is the first study to test ACT in an out-patient population with chronic medication-resistant psychotic symptoms. Although trial rigour was high, randomisation was not completely successful and drop-out rates were relatively high. The absence of a treatment-as-usual comparison means that the extent to which the observed main effect of time on many measures is attributable to specific effects of each therapy or to other factors cannot be determined, although some effectiveness of the befriending condition has been previously reported.^{12,16,17}

Befriending

Although befriending was developed originally as a control condition, the present results and previous findings^{15–17} suggest that befriending may warrant further research attention and analysis. This population is highly stigmatised, even at times in treatment settings,⁴³ and the opportunity to talk to healthcare providers about matters other than symptoms may itself be helpful to some.

Summary and directions for future research

In conclusion, objectively measured benefits of a brief ACT intervention for medication-resistant psychotic symptoms showed a moderate effect for hallucinations above those of an active control, without an effect for overall mental state or delusions. There were medium effects in favour of ACT on satisfaction and self-reported symptom benefits. The changes in process measures observed by other studies were not found here. Symptom-specific therapy refinements, improved investigation of process and attention to cognitive functioning and dose are warranted in future research.

Funding

Funded by the National Health and Medical Research Council of Australia (Project Grant 487335). We acknowledge the support of: Alfred Health, Austin Health, Eastern Health, Healthscope, Melbourne Health, Mercy Health, MIND Australia, Monash Health, NEAMI, Peninsula Health and St Vincent's Health.

Acknowledgements

This trial was undertaken through La Trobe University. We acknowledge the (non-financial) support of: Alfred Health, Austin Health, Eastern Health, Healthscope, Melbourne Health (NorthWestern Mental Health), Mercy Health, MIND Australia, Monash Health, NEAMI, Peninsula Health and St Vincent's Health. We express our gratitude to the participants.

Frances Shawyer, PhD, MAPS, MCCLP, School of Psychology and Public Health, La Trobe University, Victoria and Department of Psychiatry, School of Clinical Sciences at Monash Health, Monash University, Clayton, Victoria, Australia; **John Farhall**, PhD, FAPS, School of Psychology and Public Health, La Trobe University, Victoria and NorthWestern Mental Health, Royal Melbourne Hospital, Victoria, Australia; **Neil Thomas**, DClinPsy, School of Psychology and Public Health, La Trobe University, Victoria, School of Health Sciences, Swinburne University, Hawthorn, Victoria and Monash Alfred Psychiatry Research Centre, Melbourne, Victoria, Australia; **Steven C. Hayes**, PhD, Department of Psychology, University of Nevada, Reno, Nevada, USA; **Robert Gallop**, PhD, Department of Mathematics, Applied Statistics Program, West Chester University, West Chester, Pennsylvania, USA; **David Copolov**, PhD, MBBS, Office of the Vice-Chancellor and Discipline of Psychiatry, Monash University, Clayton, Victoria, Department of Psychiatry, University of Melbourne, Parkville, Victoria and Florey Institute of Neuroscience and Mental Health, Parkville, Victoria, Australia; **David J. Castle**, MD, FRCPsych, FRANZCP, Department of Psychiatry, University of Melbourne, Parkville, Victoria and St Vincent's Hospital Mental Health, Fitzroy, Victoria, Australia

Correspondence: John Farhall, School of Psychology and Public Health, La Trobe University, Victoria 3086, Australia. Email: j.farhall@latrobe.edu.au

First received 10 Feb 2016, final revision 9 Aug 2016, accepted 11 Aug 2016

References

- Thomas N. What's really wrong with cognitive behavioural therapy for psychosis? *Front Psychol* 2015; **6**: 323.
- Khouri B, Lecomte T, Fortin G, Masse M, Therien P, Bouchard V, et al. Mindfulness-based therapy: a comprehensive meta-analysis. *Clin Psychol Rev* 2013; **33**: 763–71.
- Hayes SC, Strosahl KD, Wilson KG. *Acceptance and Commitment Therapy: An Experiential Approach to Behavior Change*. Guilford Press, 1999.
- Hayes SC, Strosahl KD, Wilson KG. *Acceptance and Commitment Therapy: An Experiential Approach to Behavior Change (2nd edn)*. Guilford Press, 2013.
- Hayes SC. Acceptance and commitment therapy, relational frame theory and the third wave of behavior therapy. *Behav Ther* 2004; **35**: 639–65.
- Bach P, Hayes SC. The use of acceptance and commitment therapy to prevent rehospitalization of psychotic patients: a randomized controlled trial. *J Consult and Clin Psychol* 2002; **70**: 1129–39.
- Bach P, Gaudiano BA, Hayes SC, Herbert JD. Acceptance and commitment therapy for psychosis: intent to treat, hospitalization outcome and mediation by believability. *Psychosis* 2013; **5**: 166–74.
- Gaudiano BA, Herbert JD. Acute treatment of inpatients with psychotic symptoms using acceptance and commitment therapy: pilot results. *Behav Ther* 2006; **44**: 415–37.
- Thomas N, Shawyer F, Castle DJ, Copolov D, Hayes SC, Farhall J. A randomised controlled trial of acceptance and commitment therapy (ACT) for psychosis: study protocol. *BMC Psychiatry* 2014; **14**: 198.
- Kay SR. *Positive and Negative Syndromes in Schizophrenia*. Brunner/Mazel, 1991.
- Wechsler D. *Wechsler Test of Adult Reading*. Psychological Corporation, 2001.
- Shawyer F, Farhall J, Mackinnon A, Trauer, T, Sims, E, Ratcliff, K, et al. A randomised controlled trial of acceptance-based cognitive behavioural therapy for command hallucinations in psychotic disorders. *Behav Ther* 2012; **50**: 110–21.
- Bach PA. ACT with the seriously mentally ill. In *A Practical Guide to Acceptance and Commitment Therapy* (eds SC Hayes, KD Strosahl). Springer, 2005.
- Thomas N, Morris EMJ, Shawyer F, Farhall J. Acceptance and commitment therapy for voices. In *Acceptance and Commitment Therapy and Mindfulness for Psychosis* (eds EMJ Morris, LC Johns, JE Oliver): 95–111. Wiley-Blackwell, 2013.
- Bendall S, Killackey E, Jackson HJ, Gleeson J. *Befriending Manual*. ORYGEN Research Centre, University of Melbourne, 2003.
- Jackson HJ, McGorry PD, Killackey E, Bendall S, Allott K, Dudgeon P, et al. Acute-phase and 1-year follow-up results of a randomized controlled trial of CBT versus befriending for first-episode psychosis: the ACE project. *Psychol Med* 2008; **38**: 725–35.
- Sensky T, Turkington D, Kingdon D, Scott JL, Scott J, Siddle R, et al. A randomized controlled trial of cognitive-behavioural therapy for persistent symptoms in schizophrenia resistant to medication. *Arch Gen Psychiatry* 2000; **57**: 165–72.
- Bendall S, Jackson HJ, Killackey E, Allott K, Johnson T, Harrigan S, et al. The credibility and acceptability of befriending as a control therapy in a randomized controlled trial of cognitive behaviour therapy for acute first episode psychosis. *Behav Cogn Psychother* 2006; **34**: 277–91.
- Pollard S. *Measuring Treatment Fidelity in Acceptance and Commitment Therapy for Psychosis* (unpublished Master of Clinical Psychology thesis). School of Psychological Science, La Trobe University, Melbourne Australia, 2010.
- First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV-TR Axis I Disorders—Patient Edition (SCID-I/P, 1/2007 revision)*. Biometrics Research Department, New York State Psychiatric Institute, 2007.
- Haddock G, McCarron J, Tarrier N, Faragher EB. Scales to measure dimensions of hallucinations and delusions: the psychotic rating scale (PSYRATS). *Psychol Med* 1999; **29**: 879–89.
- Peters E, Joseph S, Day S, Garety P. Measuring delusional ideation: the 21-item Peters et al. Delusions Inventory (PDI). *Schizophr Bull* 2004; **30**: 1005–22.
- Birchwood M, Smith J, Cochrane R, Wetton S, Copestake S. The Social Functioning Scale. The development and validation of a new scale of social adjustment for use in family intervention programmes with schizophrenic patients. *Br J Psychiatry* 1990; **157**: 853–9.
- Hayes SC, Strosahl KD, Wilson KG, Bissett RC, Pistorello J, Toarmino D, et al. Measuring experiential avoidance: a preliminary test of a working model. *Psychol Rec* 2004; **54**: 553–78.
- Shawyer F, Ratcliff R, Mackinnon A, Farhall J, Hayes SC, Copolov D. The Voices Acceptance and Action Scale (VAAS): pilot data. *J Clin Psychol* 2007; **63**: 593–606.
- Drayton M, Birchwood M, Trower P. Early attachment experience and recovery from psychosis. *Br J Clin Psychol* 1998; **37**: 269–84.
- Attkisson CC, Zwick R. The Client Satisfaction Questionnaire: psychometric properties and correlations with service utilization and psychotherapy outcome. *Eval Prog Plan* 1982; **5**: 233–7.
- Rosenbaum PR, Rubin DB. Reducing bias in observational studies using subclassification on propensity scores. *J Am Stat Assoc* 1984; **79**: 516–24.
- Leon AC, Hedeker D. A mixed-effects quintiles-stratified propensity adjustment for effectiveness analysis of ordered categorical doses. *Stat Med* 2005; **24**: 647–58.
- Scott SC, Goldberg M, Mayo NE. Statistical assessment of ordinal outcomes in comparative studies. *J Clin Epidemiol* 1997; **50**: 45–55.
- Leucht S. Measurements of response, remission and recovery in schizophrenia and examples for their clinical application. *J Clin Psychiatry* 2014; **75** (suppl 1): 8–14.
- Feingold A. Effect sizes for growth-modeling analysis for controlled clinical trials in the same metric as for classical analysis. *Psychol Methods* 2009; **14**: 43–53.
- Burns AM, Erickson DH, Brenner CA. Cognitive-behavioral therapy for medication-resistant psychosis: a meta-analytic review. *Psychiatr Serv* 2014; **65**: 874–80.
- van der Gaag M, Valmaggia LR, Smit F. The effects of individually tailored formulation-based cognitive behavioural therapy in auditory hallucinations and delusions: a meta-analysis. *Schizophr Res* 2014; **156**: 30–7.
- Turkington D, Dudley R, Warman DM, Beck AT. Cognitive-behavioural therapy for schizophrenia: a review. *J Psychiatr Pract* 2004; **10**: 5–16.
- Birchwood M, Michail M, Meaden A, Tarrier, N, Lewis, S, Wykes, T, et al. Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. *Lancet Psychiatry* 2014; **1**: 23–33.
- Thomas N, Hayward M, Peters E, van der Gaag, M, Bentall, RP, Jenner J, et al. Psychological therapies for auditory hallucinations (voices): current status and key directions for future research. *Schizophr Bull* 2014; **40** (suppl 4): S202–12.
- Trower P, Birchwood M, Meaden A, Byrne S, Nelson A, Ross K. Cognitive therapy for command hallucinations: randomised controlled trial. *Br J Psychiatry* 2004; **184**: 312–20.
- Hooper N, Larsson A. *The Research Journey of Acceptance and Commitment Therapy (ACT)*. Palgrave Macmillan, 2015.
- Rollinson R, Smith B, Steel C, Fowler, D, Steel, C, Jolley, S, et al. Measuring adherence in CBT for psychosis: a psychometric analysis of an adherence scale. *Behav Cogn Psychother* 2008; **36**: 163–78.
- Bacon T, Farhall J, Fossey E. The active therapeutic processes of acceptance and commitment therapy for persistent symptoms of psychosis: clients' perspectives. *Behav Cogn Psychother* 2014; **42**: 402–20.
- White R, Gumley A, McTaggart J, Rattrie L, McConville D, Cleare S, et al. A feasibility study of acceptance and commitment therapy for emotional dysfunction following psychosis. *Behav Res Ther* 2011; **49**: 901–7.
- Thornicroft G, Rose D, Kassam A. Discrimination in health care against people with mental illness. *Int Rev Psychiatry* 2007; **19**: 113–22.

Online supplement DS1: Fidelity rating

Development of the ACT for Psychosis Adherence and Competence Scale - APACS

Scale Development. In the absence of any published ACT for psychosis fidelity scale, we developed a 6-item scale, the ACT for Psychosis Adherence and Competence Scale – APACS.¹⁹ The Competence subscale is not reported due to low inter-rater reliability. Only the Adherence scale was used. Adherence is measured on a 5-point (0-4) scale where ratings reflect a composite of the presence and frequency across a 50-minute session of the six ACT processes (Acceptance (Control as the problem); Defusion; Mindfulness; Self as context; Values; Willingness), as defined in our unpublished therapy manual.⁴⁴

The APACS Adherence Rating Scale:

0= the concept was not addressed

1=a little: the concept was addressed 1-2 times and briefly (up to 2 minutes spent each time) **OR** the concept was addressed once for less than 5 minutes

2=somewhat: the concept was addressed 3-5 times briefly **OR** it was addressed by the therapist once for approximately 5-14 minutes

3=considerably: the concept was addressed briefly six or more times throughout the session **OR**, a substantial amount of time was spent addressing the concept (15-25 minutes)

4= frequently: the majority of the session was spent addressing the concept

The primary score was the total of the 6 items, with the maximum realistically possible score of approximately 10. (The possible permutations of duration and frequency when summed across each of the 6 components enable a theoretical maximum score of about 14, however this is clinically implausible).

Inter-rater reliability. Twenty ACT sessions were rated by the primary rater and an independent rater agreeing on the presence of a therapy component in a session 89% of the time (107/ 120 ratings; kappa = .78, good agreement). Kappas for adherence to each

component of therapy were *Strong* or *Moderate* for each component except Defusion, where agreement was *Fair* (Table DS1).

Table DS1 Inter-rater kappa values by domain on APACS

Item	Total N	Rater 1 (M)	Rater 2 (M)	Kappa
Acceptance	20	1.20	0.95	.54** M
Defusion	20	1.05	0.70	.39** F
Mindfulness	20	2.20	1.95	.53** M
Self as context	20	0.30	0.25	.73** S
Values	20	1.30	1.15	.69** S
Willingness	20	0.55	0.25	.61** S

Note. **= $p \leq .005$; *= $p \leq .05$; ICC (C)= Intraclass correlation - consistency; S= substantial agreement; M=moderate agreement; F= fair agreement; Sl=slight agreement

Validity assessment. The first 23 participants who were enrolled in the *Lifengage* trial and who gave consent for their therapy sessions to be audio recorded were selected for inclusion in the first round of fidelity assessment (representing all available data to that date). Twelve participants were in the ACT condition and 11 in the Befriending condition. Measures used were the APACS, The Befriending Treatment Integrity Measure (BTIM – Bendall, Killackey, Jackson, & Gleeson, 2003),¹⁵ a CBT for psychosis fidelity measure the Cognitive Therapy for Psychosis Adherence Scale (R-CTPAS)⁴⁰ and the ACT Session Record, a checklist of therapy components included in a session, completed after each session by trial therapists.

Table DS2 shows descriptive data from the ACT sessions from the primary rater. The mean total APACS Adherence score of 6.54 indicates that on average, most of the session time was ratable as ACT therapy, or that therapy components were frequently identified by raters. Conversely, low mean ratings of the ACT sessions on the R-CTPAS, $M(SD)=4.17(3.29)$ and its low correlation with the APACS ($r=.25$), and low correlation ($r=.12$) with the BTIM suggested discriminant validity of the scale. Convergent validity of the APACS items was

suggested by moderate or better correlations of the presence of the 6 ACT components (Kappas from .56 – 1.0) as rated on the APACS with therapists' self-reported delivery of each ACT process recorded following each session on a checklist (the ACT Session Record).

Table DS2 Mean (SD) adherence ratings and percent of sessions with a scale item endorsed, for the APACS and validation measures using 24 randomly selected ACT sessions

Scale/Item	<i>M</i>	<i>SD</i>	Presence (% of sessions)
APACS			
Adherence Total	6.54	1.98	N/A
Acceptance	1.25	1.51	46
Defusion	0.88	1.30	33
Mindfulness	2.13	1.30	83
Self as context	0.42	1.14	12
Values	1.42	1.64	50
Willingness	0.46	1.02	21
ACT Session Record	2.45	1.01	100
R-CTPAS	4.17	3.29	79
BTIM	2.29	1.55	N/A

Note. Two sessions selected randomly from the first 12 consenting ACT participants

Fidelity analysis for the full sample

Sample. Following completion of the study a second round of fidelity rating was performed. ACT and Befriending sessions (24 sessions each) were randomly sampled from the participants who were not included in Round 1 (38 participants contributing 1-4 sessions each). The original rater was re-trained to criterion and rated all newly selected audio files. Descriptive results for ratings of ACT sessions were compared for Rounds 1 & 2 to check for any differences. Comparison of mean total adherence ratings and the 6 component means across the two rounds showed no significant differences apart from a statistically significant increase in the mean Willingness score (possibly indicating an increase in skill of therapists with practice). All sessions from each round (Round 1 ACT=24; BF=22; Round 2 ACT=24 BF=24) were pooled for the final analysis of Adherence ratings.

Adherence Results The masked rater correctly assigned to condition all ACT sessions and all befriending sessions. The mean total APACS score across all 48 ACT sessions sampled was $M(SD) = 7.06 (1.79)$ indicating on average the presence of ratable ACT processes for a majority of the session. The most frequently identified ACT components were Mindfulness and Values, being present in 43 (89.6%) and 26 (54.2%) of sampled ACT sessions respectively. Self-as-context was the least frequently rated component (3(6.3%) sessions). One sampled befriending session had a rated ACT component for values, and no ACT sessions had a befriending component rated.

Additional reference

44. Shawyer, 2008, *ACT for Psychosis Eight Session Program. Guidelines for Therapists*. School of Psychological Science, La Trobe University, Melbourne Australia, 2008.

Table DS3 Untransformed means and standard deviations for primary and secondary measures

Measure	ACT						Befriending					
	Baseline		Post-therapy		Follow-up		Baseline		Post-therapy		Follow-up	
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>
PANSS Total	49	79.2 (14.1)	43	74.6 (16.6)	41	72.5 (16.0)	47	78.6 (14.9)	42	72.6 (14.7)	36	73.9 (15.5)
PANSS Positive	49	22.8 (5.5)	43	20.6 (5.6)	41	19.3 (6.3)	47	20.8 (5.2)	42	19.1 (5.5)	36	19.6 (5.3)
PANSS Negative	49	16.9 (4.4)	43	16.8 (5.1)	41	17.2 (4.7)	47	19.2 (5.2)	42	17.8 (5.0)	36	17.8 (4.7)
PANSS General	49	39.5 (8.1)	43	37.2 (9.4)	41	35.8 (8.3)	47	38.6 (8.0)	42	35.7 (8.7)	36	36.1 (8.4)
PSYRATS AH												
Preoccupation	37	2.41 (1.32)	32	2.25 (1.05)	31	2.19 (1.49)	30	2.27 (1.41)	29	2.03 (1.59)	23	2.35 (1.50)
Origin Beliefs	36	2.50 (1.18)	31	2.16 (1.32)	30	2.53 (1.43)	31	2.65 (1.31)	29	2.34 (1.47)	22	2.32 (1.52)
Amt Distress	37	2.73 (1.02)	32	2.16 (1.27)	31	2.03 (1.40)	31	2.29 (1.37)	29	2.21 (1.52)	23	2.57 (1.62)
Int Distress	37	2.65 (0.95)	32	2.25 (1.22)	31	2.00 (1.44)	31	2.58 (1.36)	29	2.10 (1.45)	23	2.26 (1.54)
Disruption	36	1.86 (0.59)	30	1.60 (0.81)	31	1.26 (0.97)	31	1.65 (0.91)	29	1.17 (1.00)	23	1.43 (1.04)
PSYRATS DEL												
Preoccupation	47	2.53 (1.10)	41	2.07 (1.35)	38	1.92 (1.19)	42	2.79 (1.22)	40	2.13 (1.26)	32	2.19 (1.23)
Conviction	45	2.80 (1.34)	41	2.41 (1.47)	38	2.37 (1.60)	42	3.33 (1.03)	40	2.70 (1.36)	32	2.78 (1.39)
Amt Distress	47	2.77 (1.20)	41	2.20 (1.55)	39	2.05 (1.45)	42	2.69 (1.28)	40	2.13 (1.54)	32	2.34 (1.54)
Int Distress	47	2.83 (1.07)	41	1.95 (1.43)	38	1.89 (1.39)	42	2.71 (1.15)	40	2.05 (1.40)	32	2.06 (1.37)
Disruption	46	1.78 (0.63)	40	1.15 (0.80)	38	1.34 (0.94)	42	1.81 (0.86)	40	1.20 (0.97)	31	1.45 (0.89)
SFS	49	100.5(10.7)	43	99.7 (10.2)	41	100.3(10.4)	47	99.5 (8.5)	42	100.3(8.4)	35	99.7 (7.9)
No. Consults	47	29.7 (20.3)	-	-	38	32.5 (31.1)	47	30.2 (21.2)	-	-	32	22.3 (14.6)
Consult minutes	47	892 (672)	-	-	38	1240(1347)	47	911 (713)	-	-	32	909 (853)
AAQ-16	48	71.4 (13.0)	42	69.3 (11.4)	37	68.3 (12.4)	47	72.8 (10.8)	42	72.5 (10.7)	36	69.5 (9.1)
VAAS-9	35	27.7 (5.0)	33	28.4 (4.9)	26	29.3 (4.5)	29	29.3 (4.3)	29	28.9 (6.1)	20	30.3 (5.7)
RSQ	49	2.63 (1.15)	43	2.47 (1.18)	39	2.46 (1.07)	47	2.68 (1.20)	42	2.38 (1.10)	36	2.75 (1.16)

Notes: AAQ-16 = Acceptance and Action Questionnaire. PANSS = Positive and Negative Syndrome Scale. PSYRATS AH: Psychotic Symptom Rating Scales Auditory Hallucinations. PSYRATS DEL = PSYRATS Delusions. RSQ = Recovery Style Questionnaire. SFS = Social Functioning Scale. VAAS-9 = Voices Acceptance and Action Scale

Table DS4 Proportion of participants with clinically significant change ($\geq 25\%$) in PANSS total and subscale scores by condition from baseline to endpoint and baseline to follow-up

Time point	Total N	PANSS Positive <i>n</i> (%)	PANSS Negative <i>n</i> (%)	PANSS General <i>n</i> (%)	PANSS Total <i>n</i> (%)
Baseline to Post-therapy					
ACT	43	15 (34.9%)	14 (32.6%)	14 (32.6%)	11 (25.6%)
Befriending	42	11 (26.2%)	15 (35.7%)	17 (40.5%)	10 (23.8%)
Baseline to Follow-up					
ACT	41	21 (51.2%)	13 (31.7%)	14 (34.2%)	14 (34.2%)
Befriending	36	8 (22.2%)	15 (41.7%)	13 (36.1%)	14 (38.9%)

Notes: PANSS = Positive and Negative Syndrome Scale. Criterion for clinically significant change recommended by Leucht for treatment resistant populations. (Leucht S, Kane JM, Kissling W, Hamann J, Etschel E, Engel RR. What does the PANSS mean? *Schizophr Res.* 2005; 79: 231–238)

Table DS5 Means (95% CI) values and Group x Time Fixed Effects in Mixed-Effects Model Repeated Measures ANOVA for Process Measures

Measure	<u>ACT</u>			<u>Befriending</u>			<u>Group x Time</u>			
	Mean (95% CI)			Mean (95% CI)			<i>F</i>	<i>df</i>	<i>p</i>	<i>d</i>
	Baseline	Post-therapy	Follow-up	Baseline	Post-therapy	Follow-up				
AAQ-16	71.4 (68.1, 74.7)	70.3 (66.9, 73.8)	70.0 (66.4, 73.6)	72.5 (69.1, 75.9)	72.3 (68.1, 75.9)	69.8 (66.1, 73.5)	0.36	156.3	0.70	0.12 (-0.18, 0.43)
VAAS-9	27.8 (26.1, 29.5)	28.2 (26.4, 29.9)	28.7 (26.9, 30.5)	29.6 (27.7, 31.4)	28.7 (26.8, 30.6)	30.1 (28.1, 32.2)	0.66	103.6	0.52	0.17 (-0.22, 0.56)
RSQ	2.7 (2.3, 3.0)	2.5 (2.1, 2.8)	2.5 (2.1, 2.8)	2.6 (2.3, 3.0)	2.4 (2.0, 2.7)	2.7 (2.4, 3.1)	1.16	170.0	0.32	0.22 (-0.09, 0.53)

Note. AAQ-16 = Acceptance and Action Questionnaire-16; VAAS-9 = Voices Acceptance and Action Scale-9; RSQ = Recovery Style Questionnaire

BJPpsych

The British Journal of Psychiatry

Acceptance and commitment therapy for psychosis: randomised controlled trial

Frances Shawyer, John Farhall, Neil Thomas, Steven C. Hayes, Robert Gallop, David Copolov and David J. Castle

BJP published online December 15, 2016 Access the most recent version at DOI:
[10.1192/bjp.bp.116.182865](https://doi.org/10.1192/bjp.bp.116.182865)

Supplementary Material

Supplementary material can be found at:
<http://bjp.rcpsych.org/content/suppl/2016/12/05/bjp.bp.116.182865.DC1.html>

References

This article cites 0 articles, 0 of which you can access for free at:
<http://bjp.rcpsych.org/content/early/2016/12/01/bjp.bp.116.182865#BIBL>

Reprints/permissions

To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk

P<P

Published online 2016-12-15T00:05:11-08:00 in advance of the print journal.

You can respond to this article at

</letters/submit/bjprcpsych;bjp.bp.116.182865v1>

Downloaded from

<http://bjp.rcpsych.org/> on December 20, 2016
Published by The Royal College of Psychiatrists

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *The British Journal of Psychiatry* go to:
<http://bjp.rcpsych.org/site/subscriptions/>
